arXiv:hep-th/0410028v1 4 Oct 2004

Renormalizable quantum field theory as a limit of a quantum field model on the loop space.

Yu.P. Solovyov, V.V. Belokurov and E.T. Shavgulidze
Lomonosov Moscow State University, Russia
e-mail: belokur@rector.msu.ru

Abstract

A nonlocal generalization of quantum field theory in which momentum space is the space of continuous maps of a circle into $\mathbf{R}^4$ is proposed. Functional integrals in this theory are proved to exist. Renormalized quantum field model is obtained as a local limit of the proposed theory.

Consider a quantum field model defined in the following way. Let the momentum space of the theory be the Banach space

$$\mathcal{P} = C(S^1, \mathbf{R}^4)$$

of all continuous maps of a circle of a unit length into $\mathbf{R}^4$ with the norm

$$\|p\|_{\mathcal{P}} = \max_{\tau \in S^1} \|p(\tau)\| \,,$$

where $\| \cdot \|$ is the canonical norm in $\mathbf{R}^4$.

An arbitrary element of this space can be represented in the form

$$p(\tau) = r + \frac{1}{\sqrt{\lambda}} \xi(\tau) \,. \tag{1}$$

Here $\xi(\tau)$ satisfies the condition

$$\int_{S^1} \xi(\tau)\, d\tau \;=\; 0 \,. \tag{2}$$

We denote the space of maps ξ as $C_0(S^1, \mathbf{R}^4)$.

Consider the group G of three times differentiable diffeomorphisms

$$G = Diff_{+}^{\,3}\left(S^1\right) \,,$$

$$g \in G \quad \{g : \; S^1 \longrightarrow S^1 \,, \quad g'(\tau) > 0\} \,.$$

Define the action of the group on the space $\mathcal{P}$ in the following way:

$$gp(\tau) = p\left(g^{-1}(\tau)\right)\,\frac{1}{\sqrt{\left(g^{-1}\right)'(\tau)}}\,. \tag{3}$$

Consider the Wiener measure on the space $\mathcal{P}$ with the dispersion equal to $\sqrt{\lambda}$ and zero mathematical expectation

$$w_\lambda(dp) = \exp\left\{-\frac{\lambda}{2}\int\limits_{S^1} \|p'(\tau)\|^2\, d\tau\right\}\, dp\,, \quad \lambda > 0\,. \tag{4}$$

For any continuous bounded functional $F(p)$, that satisfies the inequality $|F(p)| \leq \frac{C}{1+\|p\|^5}$, the following equation is valid

$$\int\limits_{\mathcal{P}} F(p)\; w_\lambda(dp) = \int\limits_{\mathbf{R}^4}\;\int\limits_{C_0(S^1,\mathbf{R}^4)} F(r + \frac{1}{\sqrt{\lambda}}\xi)\; w_1(d\xi)\; dr\,.$$

From here, the local limit of the integral follows

$$\lim_{\lambda\to+\infty}\int\limits_{\mathcal{P}} F(p)\; w_\lambda(dp) = \int\limits_{\mathbf{R}^4} F(r)\, dr\,. \tag{5}$$

It is known that measures invariant with respect to the group G do not exist. However, the measure $w_\lambda(dp)$ is quasi-invariant. [1]

It transforms as

$$w_\lambda(d\,(gp)\,) = \exp\left\{\frac{\lambda}{4}\int\limits_{S^1} \mathcal{S}_g(\tau)\, \|p(\tau)\|^2\, d\tau\right\}\; w_\lambda(dp)\,, \tag{6}$$

Here $\mathcal{S}_g$ denotes the Schwarz derivative

$$\mathcal{S}_g(\tau) = \frac{g'''(\tau)}{g'(\tau)} - \frac{3}{2}\left(\frac{g''(\tau)}{g'(\tau)}\right)^2\,. \tag{7}$$

Consider the space E of all square-integrable over the Wiener measure functions $\varphi : \mathcal{P} \to \mathbf{C}$, satisfying the equation

$$\varphi(p) = \overline{\varphi(-p)}$$

for all $p \in \mathcal{P}$.

The space E is a Hilbert space over the real field $\mathbf{R}$ with the scalar product

$$(\varphi, \phi)_E = \int\limits_{\mathcal{P}} \varphi(p)\overline{\phi(p)}\; w_\lambda(dp)\,.$$

Functions φ and ϕ realize a regular unitary representation of the group G in the Hilbert space E:

$$g\,\varphi(p) \;=\; \varphi(g\,p)\;\exp\left\{\frac{\lambda}{8}\int\limits_{S^1} \mathcal{S}_g(\tau)\, \|p(\tau)\|^2\, d\tau\right\}\,. \tag{8}$$

Free action

$$\mathcal{A}_0[\varphi] = \int\limits_{\mathcal{P}} |\varphi(p)|^2 \, \Omega^2(p) \; w_\lambda(dp) \tag{9}$$

with

$$\Omega^2(p) = \left(\int\limits_{S^1} \frac{d\tau}{\|p(\tau)\|^2} \right)^{-1} + m^2 \tag{10}$$

is invariant with respect to the group G.

Note that

$$\lim_{\lambda\to+\infty} \mathcal{A}_0[\varphi] = \int\limits_{\mathbf{R}^4} |\varphi(r)|^2 \, \left(\|r\|^2 + m^2 \right) \, dr \, .$$

Thus, in the limit case we get the action of free scalar field.

Let us construct the interaction term. In the limit case ($\lambda \to +\infty$) it should give the usual interaction φ^4.

To get (quasi-) invariant expression we change p to $g\,p$ and consider group averaging over G with the quasi-invariant measure [2]

$$\mu_\beta(dg) = \exp\left\{ -\frac{\beta}{2} \int\limits_{S^1} \left[\left(\frac{g''(\tau)}{g'(\tau)} \right)' \right]^2 \, d\tau \right\} \, dg \, , \tag{11}$$

where β is an arbitrary positive parameter.

It is convenient to write the parameter β in the form $\beta = \alpha\lambda$ with an arbitrary positive α.

Taking into account the transformation rules for the field and the measure we propose the following formula for interaction

$$\mathcal{A}_1[\varphi] = \int\limits_{\mathcal{P}} \cdots \int\limits_{\mathcal{P}} \varphi(p_1)\varphi(p_2)\varphi(p_3)\varphi(p_4)$$

$$\int\limits_{S^1} \delta(p_1(\tau_1) - p_5(\tau_1)) \, \|p_1(\tau_1)\|^2 d\tau_1 \int\limits_{S^1} \frac{d\tau}{\|p_1(\tau)\|^2}$$

$$\int\limits_{S^1} \delta(p_2(\tau_2) - p_6(\tau_2)) \, \|p_2(\tau_2)\|^2 d\tau_2 \int\limits_{S^1} \frac{d\tau}{\|p_2(\tau)\|^2}$$

$$\int\limits_{S^1} \delta(p_3(\tau_3) - p_7(\tau_3)) \, \|p_3(\tau_3)\|^2 d\tau_3 \int\limits_{S^1} \frac{d\tau}{\|p_3(\tau)\|^2}$$

$$\int\limits_{S^1} \delta(p_4(\tau_4) - p_8(\tau_4)) \, \|p_4(\tau_4)\|^2 d\tau_4 \int\limits_{S^1} \frac{d\tau}{\|p_4(\tau)\|^2}$$

$$\int\limits_{S^1} \delta(p_5(\tau_5) + p_6(\tau_5) + p_7(\tau_5) + p_8(\tau_5)) \, \|p_5(\tau_5)\|^2 d\tau_5 \int\limits_{S^1} \frac{d\tau}{\|p_5(\tau)\|^2}$$

$$\int\limits_{G} \exp\left\{ \frac{\lambda}{8} \int\limits_{S^1} \mathcal{S}_{g_1}(\tau) \, \|p_1(\tau)\|^2 \, d\tau \right\} \; \mu_{\alpha\lambda}(dg_1) \; w_\lambda(dp_1)$$

$$\cdots$$

$$\int\limits_G \exp\left\{\frac{\lambda}{8}\int\limits_{S^1} \mathcal{S}_{g_8}(\tau)\,\|p_8(\tau)\|^2\,d\tau\right\}\ \mu_{\alpha\lambda}(dg_8)\ w_\lambda(dp_8)$$

We see that interaction in this model is nonlocal.

Using the following notations

$$\Delta(p_1,p_5)\equiv\int\limits_{\mathbf{S^1}}\frac{d\tau}{\|p_1(\tau)\|^2}\int\limits_{\mathbf{S^1}}\delta\left(p_1(\tau_1)-p_5(\tau_1)\right)\|p_1(\tau_1)\|^2 d\tau_1\,,$$

$$\mathcal{U}_{\alpha\lambda}(p)\equiv\int\limits_G \exp\left\{\frac{\lambda}{8}\int\limits_{\mathbf{S^1}}\mathcal{S}_g(\tau)\,\|p(\tau)\|^2\,d\tau\right\}\ \mu_{\alpha\lambda}(dg)\,,$$

we write it in a more compact form

$$\begin{gathered}\mathcal{A}_1[\varphi]=\int\limits_{\mathcal{P}}\cdots\int\limits_{\mathcal{P}}\ \varphi(p_1)\,\mathcal{U}_{\alpha\lambda}(p_1)\,\Delta(p_1,p_5)\\ \varphi(p_2)\,\mathcal{U}_{\alpha\lambda}(p_2)\,\Delta(p_2,p_6)\\ \varphi(p_3)\,\mathcal{U}_{\alpha\lambda}(p_3)\,\Delta(p_3,p_7)\\ \varphi(p_4)\,\mathcal{U}_{\alpha\lambda}(p_4)\,\Delta(p_4,p_8)\\ \Delta(p_5,-p_6-p_7-p_8)\\ \mathcal{U}_{\alpha\lambda}(p_5)\,\mathcal{U}_{\alpha\lambda}(p_6)\,\mathcal{U}_{\alpha\lambda}(p_7)\,\mathcal{U}_{\alpha\lambda}(p_8)\\ w_\lambda(dp_1)\,\cdots\,w_\lambda(dp_8)\,.\end{gathered}$$

Getting in mind the effects of renormalization we consider also the following addition to the action

$$\begin{gathered}\mathcal{A}_2[\varphi]=\int\limits_{\mathcal{P}}\int\limits_{\mathcal{P}}\int\limits_{\mathcal{P}}\int\limits_{\mathcal{P}}\ \varphi(p_1)\,\varphi(p_2)\\ \Delta\left(p_1,p_3\right)\ \Delta\left(p_2,p_4\right)\ \Delta\left(p_3,-p_4\right)\\ \mathcal{U}_{\alpha\lambda}(p_1)\ w_\lambda(dp_1)\,\mathcal{U}_{\alpha\lambda}(p_2)\ w_\lambda(dp_2)\\ \mathcal{U}_{\alpha\lambda}(p_3)\ w_\lambda(dp_3)\,\mathcal{U}_{\alpha\lambda}(p_4)\ w_\lambda(dp_4)\,.\end{gathered}$$

Now the total action is given by the equation

$$\mathcal{A}[\psi]=\mathcal{A}_0[\psi]+\kappa_1\mathcal{A}_1[\psi]+\kappa_2\mathcal{A}_2[\psi]\,.$$

The following theorem is valid

Theorem 1.

For any positive κ_1 and any $\psi\in E$ there exists the integral

$$\int\limits_E e^{-\mathcal{A}[\varphi]-i(\varphi,\psi)_E}d\varphi\,. \tag{12}$$

To prove this theorem note that for any $\varphi \in E$ the functional $\mathcal{A}_1[\varphi]$ is nonnegative. Therefore, the functional $e^{-\kappa_1 \mathcal{A}_1[\varphi]-\kappa_2\mathcal{A}_2[\varphi]-i(\varphi,\psi)_E}$ is bounded on E. Hence, to prove that the integral (12) converges it is sufficient to verify the existence of the integrals

$$\int_E \mathcal{A}_1[\psi] e^{-\mathcal{A}_0[\psi]} d\psi$$

and

$$\int_E \mathcal{A}_2[\psi] e^{-\mathcal{A}_0[\psi]} d\psi$$

[3].

It can be done with the help of several bounds including those that are given by the lemmas.

Lemma 1. *There exists* $c_1 > 0$ *such that* $\forall\ r \in \mathbf{R}^4$ *and* $\forall\ t_1, t_2 \in \mathbf{S^1}$ $(t_1 \neq t_2)$ *the following inequality is valid*

$$\int\limits_{\mathcal{P}} \delta\left(p(t_1)-r\right) \|p(t_1)\|^2 \, \|p(t_2)\|^2 \int\limits_{\mathbf{S^1}} \frac{d\tau}{\|p(\tau)\|^2}\, \mathcal{U}_{\alpha\lambda}(p)\ w_\lambda(dp) \leq c_1\left(1+r^4\right) .$$

Lemma 2. *There exists* $c_2 > 0$ *such that* $\forall\ t_1, t_2 \in \mathbf{S^1}$ $(t_1 \neq t_2)$ *the following inequality is fulfilled*

$$\int\limits_{\mathcal{P}} \left[\Omega^2(p_1)\right]^{-1} \|p(t_1)\|^2 \left(\|p(t_2)\|^2+1\right) \left(\int\limits_{\mathbf{S^1}} \frac{d\tau}{\|p(\tau)\|^2}\right)^2 \left(\mathcal{U}_{\alpha\lambda}(p)\right)^2\ w_\lambda(dp) \leq c_2\,.$$

We have already discussed that when $\lambda \to +\infty$ the set of measures w_λ converges in a weak sence to Lebesgue measure on $\mathbf{R}^4 \subset \mathcal{P}$.

The limit of $\mathcal{A}_0[\phi_\varphi]$ gives the action of free scalar field.

Similarly, when $\alpha \to +\infty\ \ \lambda \to +\infty$, the measure $\mathcal{U}_{\alpha\lambda}(p)\ w_\lambda(dp)$ converges in a weak sence to a Lebesgue measure. And for the limits of $\mathcal{A}_1[\phi_\varphi]$ and $\mathcal{A}_2[\phi_\varphi]$ we have the action of φ^4 model.

$$\lim_{\lambda\to+\infty\,,\alpha\to+\infty} \mathcal{A}_1[\phi_\varphi] = \int\limits_{\mathbf{R}^4}\int\limits_{\mathbf{R}^4}\int\limits_{\mathbf{R}^4}\int\limits_{\mathbf{R}^4} \delta\left(r_1+r_2+r_3+r_4\right)\ \varphi(r_1)\varphi(r_3)\varphi(r_3)\varphi(r_4)\, dr_1 dr_2 dr_3 dr_4\,.$$

$$\lim_{\lambda\to+\infty\,,\alpha\to+\infty} \mathcal{A}_2[\varphi] = \int\limits_{\mathbf{R}^4} |\varphi(r)|^2\, dr\,.$$

It can be proven that the local limit of our model is the renormalized φ^4 theory. That is, the following theorem is valid.

Theorem 2.

There are functions $\kappa_1(\theta,\alpha)$ *and* $\kappa_2(\theta,\alpha)$ *such that for all integer* $n \geq 0$ *the following limits exist*

$$\lim_{\lambda\to+\infty}\lim_{\alpha\to+\infty} \frac{\partial^n}{\partial\theta^n} \left.\frac{\int\limits_E e^{-\mathcal{A}_0[\psi]-\kappa_1(\theta,\alpha)\mathcal{A}_1[\psi]-\kappa_2(\theta,\alpha)\mathcal{A}_2[\psi]-i(\psi,\phi_\varphi)_E}\, d\psi}{\int\limits_E e^{-\mathcal{A}_0[\psi]-\kappa_1(\theta,\alpha)\mathcal{A}_1[\psi]-\kappa_2(\theta,\alpha)\mathcal{A}_2[\psi]}\, d\psi}\right|_{\theta=0} .$$

To prove this theorem we make a substitution

$$f(\tau) = \sqrt{\lambda}\left(\frac{g''(\tau)}{g'(\tau)}\right). \tag{13}$$

The measure μ looks like

$$\mu_{\alpha\lambda}(dg) = \exp\left\{-\frac{\alpha}{2}\int\limits_{S^1} \left(f'(\tau)\right)^2\, d\tau\right\} df = w_\alpha(df)\,. \tag{14}$$

Schwarz derivative $\mathcal{S}_g$ takes the form

$$\mathcal{S}_g(\tau) = \frac{1}{\sqrt{\lambda}}\, f'(\tau) - \frac{1}{2\lambda}\, f^2(\tau)\,. \tag{15}$$

Also, we have

$$\lim_{\lambda\to+\infty} \exp\left\{\frac{\lambda}{8}\int\limits_{S^1} \mathcal{S}_g(\tau)\, \|p(\tau)\|^2\, d\tau\right\} = \tag{16}$$

$$\exp\left\{-\frac{1}{16}\int\limits_{S^1} f^2(\tau)\, d\tau\, \|r\|^2 + \frac{1}{4}\int\limits_{S^1} f'(\tau)\ (r, \xi(\tau))\ d\tau\right\}.$$

Now, for the limit $\lim\limits_{\lambda\to+\infty,\ \alpha=const} \mathcal{A}_1$ we get

$$\lim_{\lambda\to+\infty,\ \alpha=const} \mathcal{A}_1[\varphi] = \int\limits_{\mathbf{R}^4} \cdots \int\limits_{\mathbf{R}^4} \int\limits_F \cdots \int\limits_F \int\limits_{C_0(S^1,\mathbf{R}^4)} \cdots \int\limits_{C_0(S^1,\mathbf{R}^4)} \varphi(r_1)\varphi(r_2)\varphi(r_3)\varphi(r_4)$$

$$\delta(r_1 - r_5)\, \delta(r_2 - r_6)\, \delta(r_3 - r_7)\, \delta(r_4 - r_8)\, \delta(r_5 + r_6 + r_7 + r_8)$$

$$\int \exp\left\{-\frac{1}{16}\int\limits_{S^1} f_1^2(\tau)\, d\tau\, \|r\|_1^2 + \frac{1}{4}\int\limits_{S^1} f_1'(\tau)\ (r_1, \xi_1(\tau))\ d\tau\right\}\ w_\alpha(df_1)\, w_1(d\xi_1)\, dr_1$$

$$\cdots$$

$$\int \exp\left\{-\frac{1}{16}\int\limits_{S^1} f_8^2(\tau)\, d\tau\, \|r\|_8^2 + \frac{1}{4}\int\limits_{S^1} f_8'(\tau)\ (r_8, \xi_8(\tau))\ d\tau\right\}\ w_\alpha(df_8)\, w_1(d\xi_8)\, dr_8\,.$$

Integrations over $r_5, ..., r_8$ and $\xi_1, ..., \xi_8$ result in

$$\lim_{\lambda\to+\infty,\ \alpha=const} \mathcal{A}_1[\varphi] = \int\limits_{\mathbf{R}^4} \cdots \int\limits_{\mathbf{R}^4} \varphi(r_1)\varphi(r_2)\varphi(r_3)\varphi(r_4)\ \int\limits_F \cdots \int\limits_F$$

$$\exp\left\{-\frac{1}{32}\int\limits_{S^1} \left[f_1^2(\tau) + f_5^2(\tau)\right]\, d\tau\ \|r_1\|^2\right\} \exp\left\{-\frac{1}{32}\int\limits_{S^1} \left[f_2^2(\tau) + f_6^2(\tau)\right]\, d\tau\ \|r_2\|^2\right\}$$

$$\exp\left\{-\frac{1}{32}\int\limits_{S^1} \left[f_3^2(\tau) + f_7^2(\tau)\right]\, d\tau\ \|r_3\|^2\right\} \exp\left\{-\frac{1}{32}\int\limits_{S^1} \left[f_4^2(\tau) + f_8^2(\tau)\right]\, d\tau\ \|r_4\|^2\right\}$$

$$w_\alpha(df_1)\ \cdots\ w_\alpha(df_8)\ \ \delta(r_1+r_2+r_3+r_4)\, dr_1\ \cdots\ dr_4\,.$$

Finally, evaluating functional integrals over f_i we obtain

$$\lim_{\lambda\to+\infty\,,\ \alpha=const} \mathcal{A}_1[\varphi] = \tag{17}$$

$$\int\limits_{\mathbf{R}^4} \dots \int\limits_{\mathbf{R}^4} \varphi(r_1)\, I\left(\alpha\,,\ \frac{1}{4}\|r_1\|\right) \cdots \varphi(r_4)\, I\left(\alpha\,,\ \frac{1}{4}\|r_4\|\right)$$

$$\delta(r_1+r_2+r_3+r_4)\, dr_1\ \cdots\ dr_4\,.$$

Here

$$I(\alpha, a) = \frac{a}{\sqrt{\alpha}}\, \frac{\exp\left(-\frac{a}{2\sqrt{\alpha}}\right)}{1-\exp\left(-\frac{a}{\sqrt{\alpha}}\right)}\,. \tag{18}$$

Similarly, for $\lim\limits_{\lambda\to+\infty\,,\ \alpha=const} \mathcal{A}_2$ we obtain

$$\lim_{\lambda\to+\infty\,,\ \alpha=const} \mathcal{A}_2[\varphi] = \tag{19}$$

$$\int\limits_{\mathbf{R}^4} \int\limits_{\mathbf{R}^4} \varphi(r_1)\, I\left(\alpha\,,\ \frac{1}{4}\|r_1\|\right)\ \ \varphi(r_2)\, I\left(\alpha\,,\ \frac{1}{4}\|r_2\|\right)$$

$$\delta(r_1+r_2)\, dr_1\, dr_2\,.$$

Note that the factors I decrease very fast at large $\|r\|$. Therefore, we can consider the above formulae as a regularization of φ^4 model with the regularization parameter α. It belongs to a set of regularizations that can be used for the proof of renormalizability of φ^4 model [4].

In a similar manner, it is possible to prove the existence of the functional integral in φ^4 model in 3-dimensional space-time

$$\lim_{\lambda\to+\infty} \lim_{\alpha\to+\infty}$$

$$\frac{\int\limits_E e^{-\mathcal{A}_0[\varphi]-\kappa_1\mathcal{A}_1[\varphi]-\kappa_2(\theta,\alpha)\mathcal{A}_2[\varphi]-i(\varphi,\chi)_E}\, d\varphi}{\int\limits_E e^{-\mathcal{A}_0[\varphi]-\kappa_1\mathcal{A}_1[\varphi]-\kappa_2(\theta,\alpha)\mathcal{A}_2[\varphi]}\, d\varphi}\,.$$

References

[1] Shavgulidze E.T. An example of the measure that is quasi-invariant with respect to the action of the group of diffeomorphisms of the circle. Funct. Anal. Appl. 1978. V. 12, N. 3. P. 55-60.

[2] Shavgulidze E.T. Some Properties of Quasi-Invariant Measures on Groups of Diffeomorphisms of the Circle. Russian Journ. Math. Phys. 2000. V. 7, N. 4, P. 464-472.

[3] Smolyanov and Shavgulidze. Continual integrals. - Moscow State University. 1990. - 150 p.

[4] Zavyalov O.I. Renormalized Feynman diagrams. - Moscow: Science, 1979. - 320 p.